



Contents lists available at ScienceDirect

NeuroImage

journal homepage: www.elsevier.com/locate/ynimg

Age-related differences in white matter integrity and cognitive function are related to APOE status

Lee Ryan^{a,*}, Katrin Walther^a, Barbara B. Bendlin^b, Lih-Fen Lue^c, Douglas G. Walker^c, Elizabeth L. Glisky^a

^a Evelyn F. McKnight Brain Institute, Department of Psychology, University of Arizona, Tucson, AZ, USA

^b Geriatric Research Education and Clinical Center, William S. Middleton Memorial Veterans Hospital, Madison, WI, USA

^c Banner Sun Health Research Institute, Sun City, AZ, USA

ARTICLE INFO

Article history:

Received 23 April 2010

Revised 16 July 2010

Accepted 19 August 2010

Available online xxx

Keywords:

Diffusion tensor imaging

Aging

Cognition

APOE

ABSTRACT

While an extensive literature is now available on age-related differences in white matter integrity measured by diffusion MRI, relatively little is known about the relationships between diffusion and cognitive functions in older adults. Even less is known about whether these relationships are influenced by the apolipoprotein (APOE) $\epsilon 4$ allele, despite growing evidence that $\epsilon 4$ increases cognitive impairment in older adults. The purpose of the present study was to examine these relationships in a group of community-dwelling cognitively normal older adults. Data were obtained from a sample of 126 individuals (ages 52–92) that included 32 $\epsilon 4$ heterozygotes, 6 $\epsilon 4$ homozygotes, and 88 noncarriers. Two measures of diffusion, the apparent diffusion coefficient (ADC) and fractional anisotropy (FA), were obtained from six brain regions—frontal white matter, lateral parietal white matter, the centrum semiovale, the genu and splenium of the corpus callosum, and the temporal stem white matter—and were used to predict composite scores of cognitive function in two domains, executive function and memory function. Results indicated that ADC and FA differed with increasing age in all six brain regions, and these differences were significantly greater for $\epsilon 4$ carriers compared to noncarriers. Importantly, after controlling for age, diffusion measures predicted cognitive function in a region-specific way that was also influenced by $\epsilon 4$ status. Regardless of APOE status, frontal ADC and FA independently predicted executive function scores for all participants, while temporal lobe ADC additionally predicted executive function for $\epsilon 4$ carriers but not noncarriers. Memory scores were predicted by temporal lobe ADC but not frontal diffusion for all participants, and this relationship was significantly stronger in $\epsilon 4$ carriers compared to noncarriers. Taken together, age and temporal lobe ADC accounted for a striking 53% of the variance in memory scores within the $\epsilon 4$ carrier group.

The results provide further evidence that APOE $\epsilon 4$ has a significant impact on the trajectory of age-related cognitive functioning in older adults. Possible mechanisms are discussed that could account for the associations between $\epsilon 4$, diffusion, and cognitive function, including the influence of $\epsilon 4$ on neural repair, oxidative stress, and the health of myelin-producing oligodendroglia.

© 2010 Elsevier Inc. All rights reserved.

Introduction

There is growing evidence of an association between the apolipoprotein (APOE) genotype and cognitive function in healthy older adults. The presence of the APOE $\epsilon 4$ allele has been related to poorer performance on tests of global cognitive functioning, episodic memory, and executive functioning (Nilsson et al., 2006; Adamson et al., 2008; Schultz et al., 2008), although a meta-analysis reviewing 38 cross-sectional studies suggests that these effects are rather small (Small et al., 2004). Longitudinal studies have revealed faster cognitive decline for $\epsilon 4$ carriers in the domains of memory (Mayeux

et al., 2001; Caselli et al., 2004; Blair et al., 2005; Swan et al., 2005), executive functioning (Swan et al., 2005), and global cognitive function (Bretsky et al., 2003; Caselli et al., 2007). Additionally, the magnitude of the cognitive decline increased as gene dose increased from one to two copies of the $\epsilon 4$ allele (Caselli et al., 2004, 2007; Blair et al., 2005).

The APOE $\epsilon 4$ allele also has been associated with structural and metabolic differences in many of the same brain regions affected by Alzheimer's disease (AD; for reviews, see Scarmeas and Stern, 2006; Cherbuin et al., 2007). Compared to noncarriers, $\epsilon 4$ carriers exhibit smaller volumes in the medial temporal lobe including the hippocampus proper (den Heijer et al., 2002; Lemaitre et al., 2005; Wishart et al., 2006), decreased cortical thickness in the entorhinal cortex and subiculum (Burggren et al., 2008), and an accelerated rate of atrophy in the hippocampus (Jak et al., 2007). Several PET studies have also demonstrated a pattern of reduced glucose metabolism in brain

* Corresponding author. Cognition & Neuroimaging Laboratories, Departments of Psychology and Neurology, University of Arizona, PO Box 210068, Tucson, AZ 85721-0068, USA. Fax: +1 520 626 7628.

E-mail address: ryant@email.arizona.edu (L. Ryan).

regions that are reminiscent of early AD, including regions of frontal, lateral temporal, parietal, and posterior cingulate cortices in $\epsilon 4$ carriers compared to noncarriers (Small et al., 2000; Reiman et al., 2005; Mosconi et al., 2008).

Recent studies have employed diffusion tensor imaging (DTI) as a method for detecting subtle changes in the integrity of brain tissue. DTI provides information about the degree and directionality of water diffusion in tissue and has been shown to be sensitive to microstructural changes in brain tissue, particularly in white matter (Le Bihan, 2003). Diffusion is most frequently reported as the apparent diffusion coefficient (ADC), a measure of mean diffusivity, and fractional anisotropy (FA), measuring the primary directionality of diffusion. In the presence of pathology such as stroke, demyelination, or inflammation, ADC is most likely to increase while FA values will decrease. Thus, DTI is a nonspecific but relatively sensitive marker of pathological changes in brain tissue, even in the absence of volumetric changes on MRI (Burgmans et al., 2010).

An extensive literature now exists on diffusion changes associated with normal aging. DTI studies have demonstrated widespread changes in diffusion across the adult lifespan with increases in ADC and decreases in FA in both anterior and posterior white matter regions. Many studies have reported age-related diffusion changes that are most prominent in the frontal lobe with relative sparing of the temporal, parietal, and occipital regions (Head et al., 2004; Hugenschmidt et al., 2008; Kochunov et al., 2007; Madden et al., 2007; Pfefferbaum et al., 2005; Salat et al., 2005; Sullivan and Pfefferbaum, 2006; Abe et al., 2008; Bhagat and Beaulieu, 2004; Deary et al., 2006; Furutani et al., 2005; Hsu et al., 2008; Hugenschmidt et al., 2008; Kochunov et al., 2007; Pagani et al., 2008; Shenkin et al., 2003, 2005; Smith et al., 2008; Zhang et al., 2005). This anterior–posterior gradient has been hypothesized to reflect the development of myelin, such that axons myelinating late in development (i.e., frontal white matter) are most vulnerable to age-related degeneration while axons myelinating early in life (i.e., posterior areas) remain relatively intact (Bartzokis, 2004). It should be noted, however, that age-related diffusion changes are also observed in posterior brain regions, particularly when the samples include a wide range of ages. For example, in a group of 52 healthy adults, ages 19 to 81 years, Kennedy and Raz (2009) reported age-related changes in both FA and ADC that extended through the brain, including frontal, temporal, parietal, and occipital lobe white matter regions.

Diffusion measures obtained from individuals with Alzheimer's disease (AD) and mild cognitive impairment (MCI) most consistently differentiate from age-matched controls in regions with a more posterior distribution, including in the cingulum bundle (Fellgiebel et al., 2005; Teipel et al., 2007; Zhang et al., 2007), parietal lobe and posterior cingulate (Medina et al., 2006; Firkbank et al., 2007; Huang et al., 2007; Ding et al., 2008; Rose et al., 2008), lateral and medial temporal lobe white matter (Fellgiebel et al., 2006; Huang et al., 2007; Yasmin et al., 2008; Xie et al., 2006; Stahl et al., 2007), and the splenium of the corpus callosum (Stahl et al., 2007). Findings in individuals with MCI tend to overlap with AD, particularly in posterior areas (Kantarci et al., 2001). In contrast, only a few studies have reported diffusion differences in the frontal lobe in MCI or AD compared to age-matched controls (Choi et al., 2005; Sydykova et al., 2007). Taken together, these findings have been described as support for the general notion that normal aging most prominently affects frontal regions, while MCI or AD results in additional changes in diffusion with a more posterior distribution (Smith et al., 2008).

Few studies, however, have examined diffusion in cognitively normal older adults who are at genetic risk for AD due to the presence of the APOE $\epsilon 4$ allele. Four studies exist to date, all focusing primarily on fractional anisotropy but interestingly using three different DTI analysis methods, one utilizing anatomically placed regions of interest, two taking a whole-brain voxel-based morphometric approach, and the third using a tract-based analysis method. The

studies are remarkably consistent in their findings, showing consistent reductions in FA values in the medial temporal lobe, although the specific region varied somewhat. Nierenberg et al. (2005) found reduced FA in the parahippocampal white matter in a sample of 14 $\epsilon 4$ carriers compared to age-matched noncarriers. Persson et al. (2006) reported that $\epsilon 4$ was associated with decreased FA in the splenium of the corpus callosum, anterior frontal lobe white matter, inferior temporal lobe white matter, the hippocampus, and the white matter of the cingulum. Smith et al. (2008), using tract-based spatial statistics, observed similar regions of decreased FA including the inferior temporal lobe white matter bilaterally, the splenium of the corpus callosum, the cingulum bundle, and anterior frontal lobe white matter. Bendlin et al. (in press), using methods similar to Persson et al. (2006) in a middle-aged cohort, did not find a main effect of APOE $\epsilon 4$ on FA but found that it interacted with parental family history of AD in medial temporal lobe white matter. The regions are similar to those reported in individuals with MCI and AD (Kantarci et al., 2001; Müller et al., 2005; Kantarci et al., 2005; Fellgiebel et al., 2005; Medina et al., 2006; Naggara et al., 2006; Cho et al., 2008; Huang et al., 2007). Smith et al. (2008) suggest that the alterations seen in at-risk individuals may reflect pathological changes associated with preclinical stages of AD.

In contrast to the quite extensive literature now available on age-related differences in white matter diffusivity, relatively little is known about the functional implications of these changes and their impact on cognitive abilities in older adults, and even less is known about the additional impact on individuals with the APOE $\epsilon 4$ allele. The studies to date have focused on very different cognitive domains or a single task within a domain, so that a consistent picture has not yet emerged. Associations have been observed on some occasions, but not others, between diffusion measures and executive function tasks (Charlton et al., 2006; Grieve et al., 2007; Shenkin et al., 2005; Sullivan et al., 2006). Speed of processing measures have also been shown to correlate with diffusivity in some samples of older adults (Bucur et al., 2008; Deary et al., 2006; Madden et al., 2004; Sullivan et al., 2006) but not others (Charlton et al., 2008; Grieve et al., 2007; Madden et al., 2007). Few studies have found an association between diffusion measures and memory (Bucur et al., 2008; Persson et al., 2006). In one recent study, Kennedy and Raz (2009) included 52 participants between 19 and 81 years of age and found that diffusion measures were not only associated with cognition, but that the relationship was region-specific: frontal white matter diffusivity predicted processing speed, posterior white matter predicted inhibition and task switching performance, while more central white matter predicted episodic memory performance.

While several studies have reported that diffusion measures predict global cognitive functioning or memory impairments in patients with MCI or AD (Hanyu et al., 1998; Bozzali et al., 2002; Müller et al., 2005; Fellgiebel et al., 2005; Ray et al., 2006; Huang et al., 2007), little is known about whether similar relationships exist for cognitively healthy individuals with genetic risk for AD. The only study to date examining the association between diffusion and cognition as a function of $\epsilon 4$ risk found no relationship between FA in the parahippocampal white matter and performance on a digit span task (Nierenberg et al., 2005). However, digit span is not generally affected in cognitively healthy APOE carriers (Small et al., 2004) or in individuals with preclinical AD (Bäckman et al., 2005), thus digit span might not be a particularly sensitive cognitive test to employ with this population. Smith et al. (2008) found changes in speed of processing associated with APOE status along with decreases in FA but did not report the correlation between the two measures.

The goal of the present paper is to further address the interactions between age, APOE $\epsilon 4$ status, and cognitive function. Specifically, does the presence of the $\epsilon 4$ allele moderate age-related changes in diffusion? Furthermore, does $\epsilon 4$ influence all brain regions equally, or only more posterior regions that have been associated with MCI

and AD? Based on the few existing studies described earlier, we hypothesize that the $\epsilon 4$ allele will increase age-related diffusion changes in more posterior regions including the temporal and parietal lobes, but not frontal regions. Finally, what is the relationship between diffusion measures and cognitive function, and does this relationship differ as a function of APOE status? While most studies of AD and MCI have focused on memory performance because of its importance as an early predictor of AD, the predominant cognitive domain associated with normal aging is executive function. Because frontal lobe diffusion changes have been associated with normal aging, we hypothesized that frontal diffusion may be associated with executive function across both $\epsilon 4$ carriers and noncarriers. In contrast, more posterior diffusion measures, including the inferior temporal lobe, may predict memory scores more strongly for $\epsilon 4$ carriers compared to noncarriers, since changes to posterior white matter diffusivity appear to be more specifically associated with the presence of the $\epsilon 4$ allele.

Methods

Participants

Volunteers included 126 healthy individuals between the ages of 52 and 92 years. All participants were living independently in the community and were recruited through newspaper advertisements and an existing database of older individuals who had participated in previous studies in our laboratory. Apolipoprotein E genotype was determined using DNA that was purified from the white blood cell fractions obtained from donated blood samples. DNA was amplified by polymerase chain reaction using primers and reaction conditions described previously (Hixson and Vernier, 1990). Amplified DNAs were digested with restriction endonuclease Hha1 and separated through 9% acrylamide gels. ApoE genotypes from each case were obtained from the patterns of digested fragments on the gels. Of the participants, 32 (25%) were heterozygous for the APOE $\epsilon 4$ allele, 6 participants (5%) were homozygous for $\epsilon 4$, and the remaining 88 participants (70%) were noncarriers, consistent with the distribution of $\epsilon 4$ found in the general population (Gerdes et al., 1992). Participants underwent medical health screening and were free from past or current neurological disorders, head injury with sequelae, psychiatric disorder, or a history of drug or alcohol abuse. No participant was taking medications that might interfere with cognitive function, such as anxiolytics. Hypertension was assessed through self-reported diagnosis of hypertension and/or current use of an antihypertensive drug. Participants were also screened for contraindications to MRI, including pacemakers, metal implants, and claustrophobia. Informed consent was obtained from each volunteer following procedures approved by the Human Subjects Committee at the University of Arizona.

Neuropsychological testing

Participants underwent a detailed neuropsychological assessment that included the Mini Mental State Exam (MMSE; Folstein et al., 1975) and the Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler, 1999) that included the Vocabulary subtest. Composite measures of memory and executive functioning were computed for each participant based on factor analyses described by Glisky (Glisky and Kong, 2008; Glisky et al., 2001). The memory factor score was based on the following tests: Logical Memory I recall total score, Verbal Paired Associate I, and Face Recognition I (Wechsler Memory Scale–III; Wechsler, 1997), Visual Paired Associates II (Wechsler Memory Scale–Revised; Wechsler, 1987), and Long-Delay Cued Recall from the California Verbal Learning Test (CVLT; Delis et al., 1987). The executive function factor score included the number of correct categories completed in the Wisconsin Card Sorting Test (WCST;

Hart et al., 1988), Mental Arithmetic total score (WAIS-R; Wechsler, 1981), total number of correct responses in the Controlled Oral Word Association Test (Spreen and Benton, 1977), Digit Span Backwards total score, and the Mental Control total score (WMS-III; Wechsler, 1997). For each participant, test scores were transformed into z scores based on Glisky and Kong's (2008) normative sample of 227 older adults and averaged to create two composite z scores for memory and executive function.

MRI acquisition and image processing

Magnetic resonance images were acquired on a GE 3.0 T Signa VH/I whole body echo-speed scanner equipped with an 8-channel phased array head coil (HD Signa Excite, General Electric, Milwaukee, WI). A set of 3-plane localizer images were obtained in order to align subsequent images. High-resolution T1-weighted whole-brain structural images were acquired using a 3D spoiled gradient-echo (3D SPGR) pulse sequence with a section thickness of 0.7 mm, no skip (TR/TE/TI = 5.1 ms/2 ms/500 ms; flip angle = 15°; matrix = 256 × 256; FOV = 260 × 260 mm²). T2 fluid-attenuated inversion recovery (T2-FLAIR) images were collected with fifty-eight axial images covering the whole brain (2.6 mm slice thickness, no gap, TE/TR/TI = 120 ms/11,000 ms/2250 ms, matrix = 256 × 192, FOV = 260 × 260 mm²). The T2-FLAIR images were used to obtain a general assessment of the amount of white matter hyperintensity using a clinical rating scale from 0 to 3 (Fazekas et al., 1987).

Diffusion tensor images were acquired with an echo planar imaging sequence corrected for spatial distortion (GE ASSET). Fifty-eight axial sections of 2.6 mm thickness, no gap, covering the whole brain were acquired (TE/TR = 71 ms/13,000 ms, matrix = 96 × 96, FOV = 250 × 250 mm²), resulting in isotropic voxels with a resolution of 2.6 mm³. Diffusion was measured in 25 directions with 2 averages ($B_0 = 1000$ s/mm², 2 NEX). While the scan resolution was somewhat lower than other relevant studies with in-plane resolution of 1.8 × 1.8 mm (for example, Smith et al., 2008; Kennedy and Raz, 2009), the somewhat large voxel size is similar to other recent studies (for example, Persson et al., 2006) and allowed us to obtain more b orientations in a reasonable amount of scanning time.

Diffusion images were resampled prior to downloading at 256 × 256 using sinc interpolation. Images were realigned to remove linear eddy current distortions using the Functional Software Library (FSL) package (<http://www.fmrib.ox.ac.uk/fsl>). DTI Studio Version 2.4 (Jiang et al., 2006; <https://www.dtistudio.org/>) software was used to compute a diffusion tensor for each voxel that included three eigenvalues and eigenvectors. Based on these values, fractional anisotropy (FA) maps and apparent diffusion coefficient (ADC) maps were computed for each participant. Diffusion maps and T1-weighted images from each individual were then oriented parallel to the anterior–posterior commissural line and coregistered to each other using SPM2 and left in native space (Wellcome Department of Cognitive Neurology, London, UK, <http://www.fil.ion.ucl.ac.uk/spm/>).

Region-of-interest (ROI) templates were created using MRICro software (Rorden, <http://www.mricro.com>) and included fully volumed regions of the frontal white matter, lateral parietal white matter, the centrum semiovale, the genu of the corpus callosum, the splenium of the corpus callosum, and the temporal stem white matter. ROIs for the frontal and parietal white matter, centrum semiovale, and temporal stem were created bilaterally on a representative T1 brain chosen from the sample (see Fig. 1). The frontal white matter was measured on 21 contiguous sections in the axial plane beginning two sections inferior to the anterior commissure. The ROI was placed in the center of the frontal white matter with the most posterior extent of the ROI corresponding to a plane level with the center of the genu. The centrum semiovale was measured on 11 contiguous sections in the coronal plane beginning five sections anterior to the anterior commissure and ending five sections posterior to the anterior




Fig. 1. Regions of interest superimposed on an individual MRI in native space. (A) Two axial and one coronal T1-weighted sections showing ROIs overlaid on the binarized white matter segmented images. The resulting ROIs include only voxels shared between the ROIs and fully volumed white matter. (B) Resulting ROIs placed on the apparent diffusion coefficient images at the same location as the T1 images. F = frontal white matter, G = genu of the corpus callosum, S = splenium of the corpus callosum, CS = centrum semiovale, TS = temporal stem white matter, P = superior parietal.

commissure. The ROI was placed on fully volumed white matter adjacent to the ventricles and superior to the caudate nucleus. The temporal stem white matter was measured on 23 contiguous coronal sections beginning at the section showing fully volumed anterior commissure. The parietal white matter was measured on 12 contiguous sections in the coronal plane beginning one section posterior to fully volumed inferior colliculi. The ROI was placed on the white matter adjacent to the ventricles and superior to the level of the lateral fissure.

Rather than using a standardized template for the corpus callosum, ROIs for the genu and splenium of the corpus callosum were drawn manually on the native space SPGRs for each participant because of the large variability in callosal shape and width across participants. The genu ROI was outlined on sagittal images beginning at the midsagittal section and continuing over the next 5 contiguous lateral sections in both hemispheres for a total of 10 sections. The posterior boundary was set at the most anterior point in the curve of the genu. The splenium ROI was drawn on consecutive sagittal images beginning at the midsagittal section and continuing over the next 6 lateral sections in both hemispheres for a total of 12 sections. The anterior boundary was set at the posterior commissure. Interrater reliability was determined for these two regions using a random sample of 10 brains. The intraclass correlations for the genu and splenium were 0.95 and 0.97, respectively, demonstrating high reliability for the procedure. The locations of the callosal ROIs are presented in Fig. 1.

To ensure that diffusion values were extracted only from white matter tissue and not partially volumed gray matter, T1 images were segmented into gray matter, white matter, and cerebrospinal fluid using SPM2. We applied the segmentation procedure used in optimized voxel-based morphometry (Good et al., 2001; Gaser, <http://www.dbm.neuro.uni-jena.de/vbm.html>), which uses information about the known location of the tissue types (priors) in addition to the signal intensity of voxels obtained from actual scans. The priors were created from a random sample of 100 participants in the present study. To minimize the potential of partial voluming effects, a conservative threshold was applied to the white matter map with a white matter probability of 0.9 to create a binarized white matter mask. The binarized mask was transformed back to native space by applying the inversion of the normalization parameters for a given individual. The ROI templates described earlier were then applied to individual participant's diffusion maps and were then multiplied by the individualized white matter mask, thereby ensuring that only fully volumed white matter was included in the ROI for a given individual. This segmentation procedure had the added benefit of removing regions of white matter hyperintensity that were visible on T2 FLAIR images. The low signal values of abnormal white matter on T1 images ensured that these regions were excluded from the white matter mask, as evident from visual comparison between the white matter mask and T2 FLAIR images. Thus, diffusion measures were only obtained from normal-appearing white matter. Diffusion values from each ROI were averaged across all extracted voxels.

Results

Demographics

Noncarrier, heterozygote, and homozygote participants are characterized in Table 1. The majority of the sample (89%) was Caucasian. The three groups were well matched on demographic and cognitive measures. One-way ANOVAs indicated no significant differences in age, years of education, cognitive functioning (MMSE, memory factor, executive factor), or clinical rating of white matter hyperintensities; all $F_s(2,123) < 1.33$, ns. Rates of hypertension were similar for noncarriers (28%) and carriers (total 29%; 31% within the heterozygote group and 17% within the homozygote group). Three individuals, all within the noncarrier group, had a diagnosis of diabetes.

For the purpose of analysing the diffusion and cognitive measures, heterozygote and homozygote groups were combined due to the small number of homozygous individuals in the sample. In addition, because of the high correlations between right and left hemisphere diffusion measures and the lack of a priori hypotheses regarding hemispheric differences, regional measures of FA and ADC were averaged across hemispheres. Other studies have also collapsed diffusion values across left and right hemispheres finding no hemisphere-specific differences in FA or ADC (for example, Kennedy and Raz, 2009). Diffusion measures were analysed using multivariate omnibus tests to control for experiment-wise error, followed by univariate tests where appropriate (detailed in sections below).

Diffusion measure omnibus tests

Omnibus tests were carried out for ADC and FA separately using a multivariate general linear model (GLM) that included the values for six white matter regions (centrum semiovale, genu, splenium, frontal lobe, parietal lobe, and temporal stem) and two between-subjects factors: age (continuous variable) and APOE $\epsilon 4$ status (carriers, noncarriers).

The omnibus test for FA showed a main effect of age, $F(1,122) = 12.51$, $p < .001$, demonstrating the previously well documented decrease in FA with increasing age. Interestingly, age did not interact with region, suggesting that age-related FA differences were generally similar across all six regions. Across all regions, FA values were lower for the $\epsilon 4$ carriers compared to noncarriers, indicated by a main effect of APOE status, $F(5,118) = 2.79$, $p < .05$. Importantly, a three-way interaction between region, age, and APOE status was obtained, $F(5,118) = 3.78$, $p < .01$, suggesting that APOE status moderated age-related differences in some but not other regions.

The omnibus test for ADC showed that this measure increased with age in all regions, indicated by the main effect of age, $F(1,122) = 91.57$, $p < .0001$, and this effect did not interact with region. In contrast to FA, however, no main effect of APOE status was observed.

Table 1

Demographics (mean \pm standard deviation) and cognitive function scores for APOE $\epsilon 4$ allele noncarrier, heterozygote, and homozygote individuals. Mean hyperintensity ratings are based on a scale of 0 to 4.

	Noncarriers	Heterozygotes	Homozygotes
Number of participants	88	32	6
Age (years)	71.6 \pm 8.8	69.2 \pm 9.7	67.0 \pm 12.2
Gender (m/f)	20/68	7/25	1/5
Hypertension (%)	28	31	17
Education (years)	15.5 \pm 2.8	15.5 \pm 2.8	15.0 \pm 2.4
MMSE	28.8 \pm 1.3	28.9 \pm 1.5	28.7 \pm 1.2
Vocabulary (raw score)	69.7 \pm 6.4	68.9 \pm 6.0	69.2 \pm 3.4
Executive factor (z score)	0.03 \pm 0.64	0.08 \pm 0.66	0.09 \pm 0.41
Memory factor (z score)	0.28 \pm 0.63	0.19 \pm 0.84	0.02 \pm 1.06
Mean hyperintensity rating	1.47 \pm 0.98	1.23 \pm 1.02	1.58 \pm 1.43

Instead, age and APOE status interacted significantly, $F(1,122) = 4.14$, $p < .05$, but did not further interact with region, suggesting that APOE status moderated age-related differences in ADC similarly across all six brain regions.

Diffusion measure univariate tests

To further elucidate the interactions between age and APOE status obtained from the omnibus tests, univariate GLM analyses were carried out separately for ADC and FA in each brain region. Although the omnibus test for ADC suggested that this step was not necessary as the APOE by age interaction applied to all regions, the univariate tests provided an estimate of the variance explained (R^2) and the related p value for each region. The univariate interaction terms and R^2 values are provided in Table 2. As expected, the univariate tests indicated that age-related increases in ADC were significantly greater for the $\epsilon 4$ carriers than noncarriers in all brain regions. For FA values, $\epsilon 4$ carriers showed significantly greater age-related decreases only within the frontal lobe and temporal lobe white matter compared to noncarriers. The genu also showed a significant interaction between age and APOE status for FA, but the interaction explained a small percentage of variance (5%) compared to the frontal and temporal lobe regions (22% and 17%, respectively). The univariate tests also showed a main effect of APOE status for FA in two regions, the frontal lobe white matter, $F(1,122) = 6.20$, $p < .05$, and the splenium, $F(1,122) = 4.76$, $p < .05$, indicating overall lower average FA values for $\epsilon 4$ carriers compared to noncarriers in these two regions.

Pearson correlations are listed in Table 3, showing the simple linear correlations between age and diffusion measures for $\epsilon 4$ carriers and noncarriers separately. Consistent with the GLM analyses described above, the table demonstrates the general increase in correlation between ADC and age across all regions, and the increasingly negative correlation between FA and age in frontal, temporal stem, and genu white matter regions for APOE $\epsilon 4$ -positive individuals compared to noncarriers.

Figs. 2 and 3 depict the simple correlations between age and ADC (Fig. 2) and age and FA (Fig. 3) in all regions. While the statistical tests were carried out comparing noncarriers to a combined group of heterozygote and homozygote individuals, the slopes of the correlation lines for the three groups are provided separately. Consistent with the GLM analyses described earlier, the dominant pattern across all brain regions appears to be an interaction between age and APOE status, such that age-related changes in diffusion values increase in the presence of the $\epsilon 4$ allele. Importantly, as depicted in Figs. 2 and 3, the increase in correlation is not due solely to the inclusion of the homozygote group but applies to those individuals with one copy of the $\epsilon 4$ allele as well.

Table 2

Univariate GLM interaction terms (F) between age and APOE status, and the percentage of variance explained (R^2) by the interaction term for each brain region.

	F	p	R^2
ADC			
Frontal	21.32	.0001	26%
Centrum	25.24	.0001	29%
Genu	22.58	.0001	27%
Temporal	30.22	.0001	33%
Parietal	19.15	.0001	24%
Splenium	17.78	.0001	22%
FA			
Frontal	16.63	.001	22%
Centrum	1.45	ns	
Genu	3.04	.05	5%
Temporal	12.74	.001	17%
Parietal	2.09	ns	
Splenium	2.35	ns	

Table 3

Pearson correlations between age and diffusion measures for APOE $\epsilon 4$ carriers ($\epsilon 4+$) and noncarriers ($\epsilon 4-$).

	ADC		FA	
	APOE $\epsilon 4-$	APOE $\epsilon 4+$	APOE $\epsilon 4-$	APOE $\epsilon 4+$
Frontal	0.38	0.73	-0.31	-0.71
Centrum	0.52	0.58	-0.07	-0.35
Genu	0.44	0.65	-0.22	-0.01
Temporal	0.42	0.78	-0.30	-0.59
Parietal	0.45	0.55	0.18	0.14
Splenium	0.38	0.67	-0.28	0.07

APOE status, diffusion, and cognition

Cognitive scores were predicted from diffusion measures from two brain regions that have been most closely associated with executive function and memory, namely the frontal white matter and temporal lobe white matter, respectively. Table 4 lists the Pearson linear correlations between the two cognitive measures and frontal ADC, frontal FA, temporal stem ADC, and temporal stem FA. Diffusion measures in both regions were consistently correlated with both memory function and executive function for the $\epsilon 4$ carriers. For the noncarriers, only frontal and temporal stem ADC measures correlated with memory scores and did not predict executive function scores.

In the present study, age was correlated with the ADC and FA measures in both groups (described earlier), and age was also correlated with cognitive function (age and memory functioning: noncarrier $r = -.39$, $\epsilon 4$ carrier $r = -.70$; age and executive functioning: noncarriers $r = -.04$, $\epsilon 4$ carriers $r = -.47$). One way to interpret the results is to consider the noncarrier group as reflecting the age-related correlations between cognition and diffusivity without the influence of APOE, which are statistically significant only between ADC and memory function (see Table 4), while the increase in correlations within the carrier group indicates the additional effect of the $\epsilon 4$ allele. To assess statistical differences between groups, the correlations were assessed with two univariate GLM models predicting memory and executive function. Predictors included the four diffusion measures as continuous variables (frontal ADC, frontal FA, temporal stem ADC, and temporal stem FA), and APOE status as a categorical variable ($\epsilon 4$ carriers, noncarriers). To control for demographic influences, gender and years of education were added as covariates along with age as a mean-centered covariate. The models tested all main effects and the interaction terms between APOE status and the four diffusion measures. Two variables, gender and temporal stem FA, did not enter into the models significantly either as main effects or as an interaction term and were therefore dropped from the analyses in order to maximize degrees of freedom. The revised model included APOE status and three diffusion measures as predictors (frontal ADC, frontal FA, and temporal stem ADC) while age and education remained as covariates.

Predicting memory scores

Results of the GLM predicting memory performance are summarized in Table 5. The model accounted for 32% of the total variance (adjusted $R^2 = .32$, $F(9,116) = 7.34$, $p < .0001$). Two important findings were observed. First, controlling for age and education, temporal stem ADC predicted memory scores while the frontal diffusion measures did not, suggesting that the relationship between memory and diffusivity is region-specific, rather than global. Second, while temporal stem ADC predicted memory significantly for both groups, APOE status interacted significantly with this factor, indicating that temporal stem ADC was a stronger predictor of memory performance for $\epsilon 4$ carriers than noncarriers, consistent with the simple correlations described earlier. The model produced a main effect of age,

$F(1,116) = 18.15$, $p < .001$, temporal stem ADC, $F(1,116) = 10.23$, $p < .001$, and a significant interaction between APOE status and temporal stem ADC, $F(1,116) = 5.59$, $p < .02$. The main effect of APOE status was marginally significant, $F(1,117) = 2.68$, $p = .10$, as was the main effect of education, $F(1,116) = 3.74$, $p = .056$. Frontal ADC and frontal FA did not approach significance either as main effects or in interactions with APOE status.

Based on these results, a post hoc model predicting memory in $\epsilon 4$ carriers alone was constructed including three predictors: age, temporal lobe ADC, and the interaction between age and temporal lobe ADC. This model predicted 53% of the variance in memory scores within the $\epsilon 4$ carrier group (adjusted $R^2 = .53$, $F(4,33) = 11.59$, $p < .001$). Importantly, the interaction term (age by temporal lobe ADC) was the only significant predictor in the model, demonstrating the importance of both variables in determining memory scores for $\epsilon 4$ carriers.

Predicting executive function scores

As summarized in Table 5, the GLM predicting executive function accounted for 22% of the variance, a significant but somewhat smaller amount of variance than was accounted for in memory performance (adjusted $R^2 = .22$, $F(9,116) = 4.95$, $p < .0001$). The model showed several interesting results that differed from the memory model. First, while controlling for age and education, executive function scores were predicted independently by both frontal ADC, $F(1,116) = 5.22$, $p < .05$, and frontal FA, $F(1,116) = 3.85$, $p < .05$, while the main effect of temporal stem ADC did not approach significance. APOE status also predicted overall executive functioning scores, indicating that $\epsilon 4$ carriers obtained overall lower executive scores than noncarriers once age was accounted for, $F(1,116) = 6.67$, $p < .01$. Finally, although the main effect of temporal stem ADC was not significant, APOE status and temporal stem ADC interacted significantly, $F(1,116) = 5.74$, $p < .05$, suggesting that temporal stem ADC values predicted executive function scores for the $\epsilon 4$ carriers, but not the noncarriers, consistent with the Pearson correlations. The main effect of years of education was also significant, $F(1,116) = 13.83$, $p < .001$ but did not interact with other variables.

Discussion

In summary, the present study demonstrated that ADC and FA diffusion measures differ with increasing age, replicating numerous prior studies. Age-related differences in ADC and FA diffusion measures were more pronounced in $\epsilon 4$ carriers than noncarriers across the whole brain, including frontal, temporal, and parietal white matter regions, as well as the corpus callosum and the centrum semiovale. Importantly, after controlling for age, region-specific diffusion measures predicted executive function and memory function, and these relationships were significantly influenced by APOE status. The results are discussed in more detail below.

Aging, APOE $\epsilon 4$, and diffusion

Three studies to date have found differences in white matter diffusion between $\epsilon 4$ carriers and noncarriers. Using a region of interest approach similar to the present study, Nierenberg et al. (2005) found lower FA as well as higher radial diffusivity in $\epsilon 4$ carriers compared to noncarriers, ages 60–75, in a region of parahippocampal white matter that was similar to the temporal stem region used in the present study. Persson et al. (2006) also found lower FA values in $\epsilon 4$ carriers compared to noncarriers in the splenium of the corpus callosum, but no differences between individuals with one or two copies of the $\epsilon 4$ allele. In addition, a voxelwise analysis revealed decreased FA in the body and splenium of the corpus callosum, the hippocampus, the occipitofrontal fasciculus, and the cingulum. Smith




Fig. 2. Scatter plots showing the relation between age and apparent diffusion coefficient (ADC) in the frontal, temporal stem, centrum semiovale, and parietal white matter as well as the genu and splenium of the corpus callosum as a function of APOE status.




Fig. 3. Scatter plots showing the relation between age and fractional anisotropy (FA) in the frontal, temporal stem, centrum semiovale, and parietal white matter as well as the genu and splenium of the corpus callosum as a function of APOE status. Note that the FA scale for the genu and splenium was changed due to the higher diffusion directionality in these regions.

Table 4

Pearson correlations between frontal and temporal stem diffusion measures and cognitive function for noncarriers ($\epsilon 4-$) and carriers ($\epsilon 4+$) of the APOE $\epsilon 4$ allele.

	Memory functioning		Executive functioning	
	APOE $\epsilon 4-$	APOE $\epsilon 4+$	APOE $\epsilon 4-$	APOE $\epsilon 4+$
Frontal ADC	-.18*	-.49*	.16	-.42*
Frontal FA	.14	.43*	.04	.46*
Temporal stem ADC	-.26*	-.65*	.11	-.56*
Temporal stem FA	.14	.57*	.04	.50*

* $p < .05$, two-tailed.

et al. (2008), using whole-brain spatial statistics analysis methods, compared two groups of women ranging in age from 40 to 90 years; 42 women were $\epsilon 4$ -positive with a family history of Alzheimer's disease and 23 women were noncarriers. Similar to the results from Persson et al. (2006), they found lower FA values in the inferior temporal lobe, splenium, anterior cingulum, and frontal white matter. Common across all three studies, despite large methodological differences, was the finding of lower FA values in the parahippocampal/inferior temporal lobe white matter for $\epsilon 4$ carriers relative to noncarriers.

While our results are generally consistent with previous literature, the present study is unique in several ways. First, the three studies described above focused primarily on FA (radial diffusion was considered in Nierenberg et al., 2005), while the present study evaluated both FA and ADC. These two measures tend to be negatively correlated to a moderate degree within white matter tracts, but they may be sensitive to different aspects of pathological change in white matter (Song et al., 2002). We found that ADC, as well as FA, differed across all six regions of the brain sampled here as a function of age and APOE status. Second, rather than comparing mean differences across groups, the present study focused on the trajectory of age-related changes in diffusion measures. Taking this approach highlights the finding that the $\epsilon 4$ effect does not apply equally across all older adults, but rather appears to affect individuals differentially in increasing decades, as indicated by the interactions between APOE status, age, and diffusion values. The differential effect of $\epsilon 4$ over the life span may account in part for the lack of a main effect of $\epsilon 4$ on diffusion found in Bendlin et al. (in press), where participants were largely middle-aged. Although the cross-sectional design of the present study precludes making strong statements regarding longitudinal change as a function of age, the pattern of results suggests that the $\epsilon 4$ allele does not impart an additive effect on diffusion across the entire older adult age range, but rather changes the trajectory of normal aging.

Table 5

Univariate GLM models showing significant predictors of memory and executive function scores for APOE $\epsilon 4$ carriers and noncarriers.

Executive function		Adjusted $R^2 = .22, p < .0001$	
Predictor	F		p
Years education	13.83		.001
Frontal ADC	5.23		.01
Frontal FA	3.85		.05
APOE status	6.67		.01
APOE \times temporal ADC	5.74		.02
Memory function		Adjusted $R^2 = .32, p < .0001$	
Predictor	F		p
Age	18.15		.001
Temporal ADC	10.23		.002
APOE \times temporal ADC	5.59		.02

The present results suggest that age-related differences in diffusion were observed in all regions of the brain, both anterior and posterior. Previous studies have suggested that aging results in more pronounced changes in frontal white matter (e.g., Bartzokis, 2004) with relative sparing of posterior regions, although more recent studies including Kennedy and Raz (2009) reported age-related changes in both FA and ADC that extended throughout both anterior and posterior brain regions. In our study, ADC was similarly correlated with age in all six brain regions, while FA was most strongly correlated with age within the frontal lobe and the temporal stem. The results are consistent with recent work by Zhang et al. (2008), comparing the pattern of age-related differences for FA and mean diffusivity (similar to ADC) across various brain regions. They found that both measures differed with age (FA decreased while mean diffusivity increased) in frontal areas, the corpus callosum, periventricular white matter, and brain stem. However, mean diffusivity differences were more widely distributed than FA differences encompassing both anterior and posterior brain regions, and several regions showed significant mean diffusivity differences without decreases in FA (bilateral cingulum, bilateral thalamus, caudate nuclei, the centrum semiovale, and the medial temporal lobe). The relatively large sample size in the present study and separating the variance associated with both aging and APOE status likely increased the power to detect differences in all regions. The present results suggest that age-related differences in diffusion are best described as affecting the whole brain rather than frontal regions only.

Importantly, we found that $\epsilon 4$ exacerbated age-related diffusion differences in all regions of the brain that were sampled, both anterior and posterior. This is consistent with a recent study (Smith et al., 2008) showing lower FA values in the white matter of the frontal lobe, inferior temporal lobe, splenium, and anterior cingulum bundle. The results of these two studies are somewhat at odds with the majority of diffusion studies of MCI and AD patients suggesting that MCI/AD primarily affects posterior regions with relative sparing of frontal lobes. Smith et al. (2008) have suggested that the alterations observed in cognitively normal $\epsilon 4$ carriers reflect pathological changes associated with preclinical stages of AD, rather than normal aging. One might expect, then, that $\epsilon 4$ differences in diffusion should be most prominent in posterior regions rather than across the whole brain, reflecting the known distribution of early AD pathology. However, at least two studies have reported differences in frontal diffusion in MCI and AD groups compared to age-matched controls (Choi et al., 2005; Sydykova et al., 2007). The findings of this study and Smith et al. (2008) suggest that $\epsilon 4$ has widespread effects on diffusion that include both posterior regions associated with early AD pathology as well as more anterior regions of frontal white matter typically associated with normal aging. Some of the difference across studies may arise because few studies have separated participants based on APOE status. Although we did not have sufficient homozygotic individuals in the study to assess the differential effect of one versus two copies of $\epsilon 4$, the present study demonstrated that the effect of $\epsilon 4$ was not being driven solely by the homozygotic individuals. At the very least, we can say that having even a single copy of the $\epsilon 4$ allele is sufficient to increase the amount of age-related diffusion difference in the white matter regions and that the pattern of results is as strong, if not stronger, for those individuals with two copies of the $\epsilon 4$ allele.

The relation between diffusion and cognitive function

While many studies have found that diffusion measures differ with increasing age, fewer studies have focused on the association between diffusion and cognition, and even fewer on whether the association between diffusion and cognition is influenced by the presence of the $\epsilon 4$ allele. In the present study, we found that diffusion measures predicted cognitive function in a region-specific way, and that $\epsilon 4$

had a significant influence on these associations. More specifically, regardless of APOE status, frontal white matter diffusion measures (ADC and FA) independently predicted executive function scores for all participants, while temporal lobe ADC additionally predicted executive function for $\epsilon 4$ carriers, but not noncarriers. Memory scores were predicted specifically by temporal lobe ADC but not frontal diffusion measures for all participants, although the relationship between temporal lobe ADC and memory scores was significantly strengthened in $\epsilon 4$ carriers compared to noncarriers. Taken together, age and temporal lobe ADC accounted for a striking 53% of the variance in memory scores within the $\epsilon 4$ carrier group. It should be noted that ADC and FA values in the temporal lobe were highly correlated with one another. Once ADC was included in the multivariate models, FA did not add much predictive value to the model.

The results of previous studies on the influence of $\epsilon 4$ on cognitive function are mixed. Adamson et al. (2008) found no age by APOE interaction for memory function, but participants in the study were younger (ages 50–76) compared to the present study, an age range where $\epsilon 4$ had relatively limited effect on memory function. Consistent with our findings, Nilsson et al. (2006) found episodic memory decreased in $\epsilon 4$ carriers compared to noncarriers, but only in the old-old (ages 70–85), not young-old (ages 50–65), or young (ages 35–50). Somewhat inconsistent with our findings, a recent paper by Raz et al. (2009) found that $\epsilon 4$ resulted in greater age-related differences in executive function but not memory function. However, their sample included 189 individuals from ages 18 to 82 years. Such a sample with fewer participants in the older age range would be less likely to observe $\epsilon 4$ effects that may be present only in the 7th decade and beyond, as suggested by the present findings.

An important question is the extent to which the results are driven by the inclusion of individuals in preclinical stages of AD, or whether the APOE effects observed here are influencing the trajectory of normal aging. In other studies, the term “presymptomatic” has been applied to all individuals who are $\epsilon 4$ -positive but otherwise cognitively healthy (e.g., Smith et al., 2008). However, this term implies that each person who is $\epsilon 4$ positive will go on to develop AD. Recent studies suggest that $\epsilon 4$ carries the greatest AD risk for individuals between 60 and 69 years. In this age group, nearly 50% of patients diagnosed with AD are $\epsilon 4$ carriers. At age 70 and beyond, the frequency of $\epsilon 4$ allele decreases in cognitively normal populations (Sando et al., 2008; Davidson et al., 2007). In the present study, more than half of the sample was age 70 or above, and this is the group in which $\epsilon 4$ had the strongest effect on cognitive functioning. We suggest that APOE may have dual effects. In earlier years, particularly in the 5th and 6th decades, the presence of $\epsilon 4$ may be related more directly to the development of pathology that results in a diagnosis of MCI or probable AD. In later decades, the influence of $\epsilon 4$ on the trajectory of normal aging may become more clearly visible, because a significant number of individuals who will develop AD as a function of $\epsilon 4$ risk have already been removed from normal aging samples. While this question can never be completely answered without longitudinal data, the present results suggest that $\epsilon 4$ may have a negative effect on the normal development of age-related changes in executive function and memory per se that may be separate from the development of the pathology associated with early AD. We are now following the sample longitudinally to determine the rate of decline in cognition as a function of APOE status and diffusion, and whether some of these individuals go on within a 2- to 3-year period to develop MCI or AD.

What could be the underlying mechanism that accounts for the associations observed here between $\epsilon 4$, diffusion, and normal age-related changes in cognitive function? We consider two possibilities, but clearly others could exist and in fact it is likely that several of these mechanisms interact with one another to produce cognitive decline. Separate from its role in the development of beta-amyloid, we first

consider the influence of $\epsilon 4$ on the function of neurons, and second, the influence of $\epsilon 4$ on inflammation and the health of myelin-producing oligodendroglia.

APOE is critical for maintaining the health and function of neurons through its role in redistributing lipids among CNS cells for normal lipid homeostasis, neuronal repair, synapse maintenance, and scavenging toxins. For example, through its lipid transport function, APOE $\epsilon 3$ and $\epsilon 2$ are effective in remodeling and repairing synaptodendritic connections, but $\epsilon 4$ is much less effective (Mahley and Rall, 2000; reviewed in Mahley et al., 2006). The presence of $\epsilon 4$ therefore results in increased injury to cells in the presence of stressors such as oxidative stress, ischemia, excess beta-amyloid production, SOD1 mutations, inflammation, and the aging process itself. In addition, APOE $\epsilon 4$ disrupts mitochondrial regulation of energy and glucose metabolism in neurons. An association between $\epsilon 4$ genotype and altered CNS glucose metabolism has been demonstrated in both AD patients and cognitively normal older adults (Small et al., 2000; Reiman et al., 2001, 2004). Thus, even in adults without significant beta-amyloid deposition, $\epsilon 4$ is likely to result in reduced neuronal glucose utilization due to altered mitochondrial activity and increased levels of cell damage due to a variety of CNS stressors that are known to increase with normal aging.

In comparison to $\epsilon 2$ and $\epsilon 3$, APOE $\epsilon 4$ results in an imbalance between pro- and anti-inflammatory cytokines that promotes a pro-inflammatory state and increases oxidative stress (reviewed in Jofre-Monseny et al., 2008). Oligodendrocytes that are responsible for the production and maintenance of myelin are very vulnerable to oxidative stress, inflammatory cytokines, ischemia, and excitotoxic neurotransmitters (Ludwin, 1997). APOE $\epsilon 4$ could contribute to accelerated myelin breakdown that underlies age-related changes in diffusion. The death of oligodendrocytes can activate the generation of new oligodendrocytes and remyelination of the axon, but the resulting myelin sheaths are thinner with shorter internodes than the original myelin (Peters, 2002, 2009). Such alterations in myelin would result in decreased conduction velocity and disrupt the coordinated timing of neuronal signals across multiple brain regions. Damage to the myelin sheath would likely result in decreased FA diffusion and increased ADC and possibly contribute to the cognitive impairments observed in older adults.

The notion that $\epsilon 4$ is related to inflammatory stress is consistent with several findings, both from AD and normal aging. For instance, meta-analyses of epidemiological studies show a 50% or better decrease in risk for developing AD among users of nonsteroidal anti-inflammatory drugs (NSAIDs) compared to nonusers (Etminan et al., 2003; McGeer et al., 1996; Szekely et al., 2004). In contrast to the overwhelmingly positive results of epidemiological studies, clinical trials of NSAIDs for the treatment of probable AD have yielded disappointing results (Aisen et al., 2002, 2003; Reines et al., 2004; Scharf et al., 1999). These studies suggested that NSAIDs may be useful prophylactically rather than as a treatment for the disorder after the first symptoms appear. However, a large-scale primary prevention study that enrolled over 2500 participants (ADAPT Research Group et al., 2007) also yielded negative results, although the study included only individuals over the age of 70 years. Few studies have considered anti-inflammatory drugs in the context of normal aging. A recent study from our laboratory focusing on cognitively normal older adults demonstrated that NSAID drug users had significantly less age-related decreases in gray and white matter volumes compared to age-matched controls (Walther et al., 2009). These results suggest that drug treatments or life-style changes that combat inflammation may slow age-related changes in brain structure and function (Jofre-Monseny et al., 2008). This is a promising line of research that warrants further investigation.

In summary, the present study provides evidence that age-related differences in white matter diffusion interact with APOE status, which together predict cognitive functions of older adults in the areas of

executive function and memory. We suggest that the presence of $\epsilon 4$ plays an important role in the trajectory of normal aging, possibly via mechanisms related to neural repair, inflammation, and oxidative stress. Our study suggests that the combination of $\epsilon 4$ and diffusion MRI is a good predictor of who will experience cognitive impairments as they age. Future studies may determine whether $\epsilon 4$ also results in faster rates of decline and whether drug or lifestyle interventions can ameliorate these cognitive effects.

Acknowledgments

The project was supported by the Arizona Biomedical Research Commission, the Arizona Alzheimer's Research Consortium (HB 2354, Dept. of Health Services, Arizona), the Evelyn F. McKnight Brain Institute, the National Institute on Aging (P30 AG19610 Arizona Alzheimer's Disease Core Center), and the National Institute on Neurologic Disorder and Stroke (RO1 NS044107). We thank Ms. Kristina Irwin for coordinating and administering neuropsychological testing, and Dr. Thomas Beach at Banner Sun Health Research Institute for his generous support.

References

- Abe, O., Yamasue, H., Aoki, S., Suga, M., Yamada, H., Kasai, K., Masutani, Y., Kato, N., Ohtomo, K., 2008. Aging in the CNS: comparison of gray/white matter volume and diffusion tensor data. *Neurobiol. Aging* 29, 102–116.
- Adamson, M.M., Landy, K.M., Duong, S., Fox-Bosetti, S., Ashford, J.W., Murphy, G.M., Weiner, M., Taylor, J.L., 2008. Apolipoprotein E epsilon4 influences on episodic recall and brain structures in aging pilots. *Neurobiol. Aging* doi:10.1016/j.neurobiolaging.2008.07.017.
- ADAPT Research Group, Lyketsos, C.G., Breitner, J.C., Green, R.C., Martin, B.K., Meinert, C., Piantadosi, S., Sabbagh, M., 2007. Naproxen and celecoxib do not prevent AD in early results from a randomized controlled trial. *Neurology* 68, 1800–1808.
- Aisen, P.S., Schmeidler, J., Pasinetti, G.M., 2002. Randomized pilot study of nimesulide treatment in Alzheimer's disease. *Neurology* 58, 1050–1054.
- Aisen, P.S., Schafer, K.A., Grundman, M., Pfeiffer, E., Sano, M., Davis, K.L., Farlow, M.R., Jin, S., Thomas, R.G., Thal, L.J., 2003. Effects of rofecoxib or naproxen vs placebo on Alzheimer disease progression: a randomized controlled trial. *JAMA* 289, 2819–2826.
- Bäckman, L., Jones, S., Berger, A.K., Laukka, E.J., Small, B.J., 2005. Cognitive impairment in preclinical Alzheimer's disease: a meta-analysis. *Neuropsychology* 19, 520–531.
- Bartzokis, G., 2004. Age-related myelin breakdown: a developmental model of cognitive decline and Alzheimer's disease. *Neurobiol. Aging* 25, 5–18.
- Bendlin, B.B., Ries, M.L., Canu, E., Lazar, M., Alexander, A.L., Carlsson, C.M., Sager, M.A., Asthana, S., Johnson, S.C., 2010. White matter is altered with parental family history of Alzheimer's disease. *Alzheimer's Dement* 6, 394–403.
- Bhagat, Y.A., Beaulieu, C., 2004. Diffusion anisotropy in subcortical white matter and cortical gray matter: changes with aging and the role of CSF-suppression. *J. Magn. Reson. Imaging* 20, 216–227.
- Blair, C.K., Folsom, A.R., Knopman, D.S., Bray, M.S., Mosley, T.H., Boerwinkle, E., Atherosclerosis Risk in Communities (ARIC) Study Investigators, 2005. APOE genotype and cognitive decline in a middle-aged cohort. *Neurology* 64, 268–276.
- Bozzali, M., Falini, A., Franceschi, M., Cercignani, M., Zuffi, M., Scotti, G., Comi, G., Filippi, M., 2002. White matter damage in Alzheimer's disease assessed in vivo using diffusion tensor magnetic resonance imaging. *J. Neurol. Neurosurg. Psychiatry* 72, 742–746.
- Bretsky, P., Guralnik, J.M., Launer, L., Albert, M., Seeman, T.E., 2003. The role of APOE-epsilon4 in longitudinal cognitive decline: MacArthur Studies of Successful Aging. *Neurology* 60, 1077–1081.
- Bucur, B., Madden, D.J., Spaniol, J., Provenzale, J.M., Cabeza, R., White, L.E., Huettel, S.A., 2008. Age-related slowing of memory retrieval: contributions of perceptual speed and cerebral white matter integrity. *Neurobiol. Aging* 29, 1070–1079.
- Burggren, A.C., Zeineh, M.M., Ekstrom, A.D., Braskie, M.N., Thompson, P.M., Small, G.W., Bookheimer, S.Y., 2008. Reduced cortical thickness in hippocampal subregions among cognitively normal apolipoprotein E $\epsilon 4$ carriers. *Neuroimage* 41, 1177–1183.
- Burgmans, S., van Boxtel, M.P., Gronenschild, E.H., Vuurman, E.F., Hofman, P., Uylings, H.B., Jolles, J., Raz, N., 2010. Multiple indicators of age-related differences in cerebral white matter and the modifying effects of hypertension. *Neuroimage* 49, 2083–2093.
- Caselli, R.J., Reiman, E.M., Osborne, D., Hentz, J.G., Baxter, L.C., Hernandez, J.L., Alexander, G.G., 2004. Longitudinal changes in cognition and behavior in asymptomatic carriers of the APOE $\epsilon 4$ allele. *Neurology* 62, 1990–1995.
- Caselli, R.J., Reiman, E.M., Locke, D.E., Hutton, M.L., Hentz, J.G., Hoffman-Snyder, C., Woodruff, B.K., Alexander, G.E., Osborne, D., 2007. Cognitive domain decline in healthy apolipoprotein E epsilon4 homozygotes before the diagnosis of mild cognitive impairment. *Arch. Neurol.* 64, 1306–1311.
- Charlton, R.A., Barrick, T.R., McIntyre, D.J., Shen, Y., O'Sullivan, M., Howe, F.A., Clark, C.A., Morris, R.G., Markus, H.S., 2006. White matter damage on diffusion tensor imaging correlates with age-related cognitive decline. *Neurology* 66, 217–222.
- Charlton, R.A., Landau, S., Schiavone, F., Barrick, T.R., Clark, C.A., Markus, H.S., Morris, R. G., 2008. A structural equation modeling investigation of age-related variance in executive function and DTI measured white matter damage. *Neurobiol. Aging* 29, 1547–1555.
- Cherbuin, N., Leach, L.S., Christensen, H., Anstey, K.J., 2007. Neuroimaging and APOE genotype: a systematic qualitative review. *Dement. Geriatr. Cogn. Disord.* 24, 348–362.
- Cho, H., Yang, D.W., Shon, Y.M., Kim, B.S., Kim, Y.I., Choi, Y.B., Lee, K.S., Shim, Y.S., Yoon, B., Kim, W., Ahn, K.J., 2008. Abnormal integrity of corticocortical tracts in mild cognitive impairment: a diffusion tensor imaging study. *J. Korean Med. Sci.* 23, 477–483.
- Choi, S.J., Lim, K.O., Monteiro, I., Reisberg, B., 2005. Diffusion tensor imaging of frontal white matter microstructure in early Alzheimer's disease: a preliminary study. *J. Geriatr. Psychiatry Neurol.* 18, 12–19.
- Davidson, Y., Gibbons, L., Pritchard, A., Hardicre, J., Wren, J., Stopford, C., Julien, C., Thompson, J., Payton, A., Pickering-Brown, S.M., Pendleton, N., Horan, M.A., Burns, A., Purandare, N., Lendon, C.L., Neary, D., Snowden, J.S., Mann, D.M., 2007. Apolipoprotein E epsilon4 allele frequency and age at onset of Alzheimer's disease. *Dement. Geriatr. Cogn. Disord.* 23, 60–66.
- Deary, I.J., Bastin, M.E., Pattie, A., Clayden, J.D., Whalley, L.J., Starr, J.M., Wardlaw, J.M., 2006. White matter integrity and cognition in childhood and old age. *Neurology* 66, 505–512.
- Delis, D.C., Kramer, J., Kaplan, E., Ober, B.A., 1987. The California Verbal Learning Test. Psychological Corporation, San Antonio, TX.
- den Heijer, T., Oudkerk, M., Launer, L.J., van Duijn, C.M., Hofman, A., Breteler, M.M., 2002. Hippocampal, amygdalar, and global brain atrophy in different apolipoprotein E genotypes. *Neurology* 59, 746–748.
- Ding, B., Chen, K.M., Ling, H.W., Zhang, H., Chai, W.M., Li, X., Wang, T., 2008. Diffusion tensor imaging correlates with proton magnetic resonance spectroscopy in posterior cingulate region of patients with Alzheimer's disease. *Dement. Geriatr. Cogn. Disord.* 25, 218–225.
- Etmann, M., Gill, S., Samii, A., 2003. Effect of non-steroidal anti-inflammatory drugs on risk of Alzheimer's disease: systematic review and meta-analysis of observational studies. *BMJ* 327, 128.
- Fazekas, F., Chawluk, J.B., Alavi, A., Hurtig, H.I., Zimmerman, R.A., 1987. MRI signal abnormalities at 1.5 T in Alzheimer's dementia and normal aging. *AJNR* 8, 421–426.
- Fellgiebel, A., Müller, M.J., Wille, P., Dellani, P.R., Scheurich, A., Schmidt, L.G., Stoeter, P., 2005. Color-coded diffusion-tensor-imaging of posterior cingulate fiber tracts in mild cognitive impairment. *Neurobiol. Aging* 26, 1193–1198.
- Fellgiebel, A., Dellani, P.R., Greverus, D., Scheurich, A., Stoeter, P., Müller, M.J., 2006. Predicting conversion to dementia in mild cognitive impairment by volumetric and diffusivity measurements of the hippocampus. *Psychiatry Res.* 146, 283–287.
- Firbank, M.J., Blamire, A.M., Krishnan, M.S., Teodorczuk, A., English, P., Gholkar, A., Harrison, R., O'Brien, J.T., 2007. Atrophy is associated with posterior cingulate white matter disruption in dementia with Lewy bodies and Alzheimer's disease. *Neuroimage* 36, 1–7.
- Folstein, M.F., Folstein, S.E., McHugh, P.R., 1975. "Mini-mental state." A practical method for grading the cognitive state of patients for the clinician. *J. Psychiatr. Res.* 12, 189–198.
- Furutani, K., Harada, M., Minato, M., Morita, N., Nishitani, H., 2005. Regional changes of fractional anisotropy with normal aging using statistical parametric mapping (SPM). *J. Med. Invest.* 52, 186–190.
- Gerdes, L.U., Klausen, I.C., Sihm, I., Faergeman, O., 1992. Apolipoprotein E polymorphism in a Danish population compared to findings in 45 other study populations around the world. *Genet. Epidemiol.* 9, 155–167.
- Glisky, E.L., Kong, L.L., 2008. Do young and older adults rely on different processes in source memory tasks? A neuropsychological study. *J. Exp. Psychol. Learn. Mem. Cogn.* 34, 809–822.
- Glisky, E.L., Rubin, S.R., Davidson, P.S., 2001. Source memory in older adults: an encoding or retrieval problem? *J. Exp. Psychol. Learn. Mem. Cogn.* 27, 1131–1146.
- Good, C.D., Johnsrude, I.S., Ashburner, J., Henson, R.N., Friston, K.J., Frackowiak, R.S., 2001. A voxel-based morphometric study of ageing in 465 normal adult human brains. *Neuroimage* 14, 21–36.
- Grieve, S.M., Williams, L.M., Paul, R.H., Clark, C.R., Gordon, E., 2007. Cognitive aging, executive function, and fractional anisotropy: a diffusion tensor MR imaging study. *AJNR Am. J. Neuroradiol.* 28, 226–235.
- Hanyu, H., Sakurai, H., Iwamoto, T., Takasaki, M., Shindo, H., Abe, K., 1998. Diffusion-weighted MR imaging of the hippocampus and temporal white matter in Alzheimer's disease. *J. Neurol. Sci.* 156, 195–200.
- Hart, R.P., Kwentus, J.A., Wade, J.B., Taylor, J.R., 1988. Modified Wisconsin Card Sorting Test in elderly normal, depressed and demented patients. *Clin. Neuropsychol.* 2, 49–56.
- Head, D., Buckner, R.L., Shimony, J.S., Williams, L.E., Akbudak, E., Conturo, T.E., McAvoy, M., Morris, J.C., Snyder, A.Z., 2004. Differential vulnerability of anterior white matter in nondemented aging with minimal acceleration in dementia of the Alzheimer type: evidence from diffusion tensor imaging. *Cereb. Cortex* 14, 410–423.
- Hixson, J.E., Vernier, D.T., 1990. Restriction isotyping of human apolipoprotein E by gene amplification and cleavage with HhaI. *J. Lipid Res.* 31, 545–548.
- Hsu, J.L., Leemans, A., Bai, C.H., Lee, C.H., Tsai, Y.F., Chiu, H.C., Chen, W.H., 2008. Gender differences and age-related white matter changes of the human brain: a diffusion tensor imaging study. *Neuroimage* 39, 566–577.
- Huang, J., Friedland, R.P., Auchus, A.P., 2007. Diffusion tensor imaging of normal-appearing white matter in mild cognitive impairment and early Alzheimer disease:

- preliminary evidence of axonal degeneration in the temporal lobe. *AJNR Am. J. Neuroradiol.* 28, 1943–1948.
- Hugenschmidt, C.E., Peiffer, A.M., Kraft, R.A., Casanova, R., Deibler, A.R., Burdette, J.H., Malsjian, J.A., Laurienti, P.J., 2008. Relating imaging indices of white matter integrity and volume in healthy older adults. *Cereb. Cortex* 18, 433–442.
- Jak, A.J., Houston, W.S., Nagel, B.J., Corey-Bloom, J., Bondi, M.W., 2007. Differential cross-sectional and longitudinal impact of APOE genotype on hippocampal volumes in nondemented older adults. *Dement. Geriatr. Cogn. Disord.* 23, 382–389.
- Jiang, H., van Zijl, P.C., Kim, J., Pearlson, G.D., Mori, S., 2006. DtiStudio: resource program for diffusion tensor computation and fiber bundle tracking. *Comput. Meth. Programs Biomed.* 81, 106–116.
- Jofre-Monseny, L., Minihane, A.M., Rimbach, G., 2008. Impact of apoE genotype on oxidative stress, inflammation and disease risk. *Mol. Nutr. Food Res.* 52, 131–145.
- Kantarci, K., Jack Jr., C.R., Xu, Y.C., Campeau, N.G., O'Brien, P.C., Smith, G.E., Ivnik, R.J., Boeve, B.F., Kokmen, E., Tangalos, E.G., Petersen, R.C., 2001. Mild cognitive impairment and Alzheimer disease: regional diffusivity of water. *Radiology* 219, 101–107.
- Kantarci, K., Petersen, R.C., Boeve, B.F., Knopman, D.S., Weigand, S.D., O'Brien, P.C., Shiung, M.M., Smith, G.E., Ivnik, R.J., Tangalos, E.G., Jack Jr., C.R., 2005. DWI predicts future progression to Alzheimer disease in amnesic mild cognitive impairment. *Neurology* 64, 902–904.
- Kennedy, K.M., Raz, N., 2009. Aging white matter and cognition: differential effects of regional variations in diffusion properties on memory, executive functions, and speed. *Neuropsychologia* 47, 916–927.
- Kochunov, P., Thompson, P.M., Lancaster, J.L., Bartzokis, G., Smith, S., Coyle, T., Royall, D. R., Laird, A., Fox, P.T., 2007. Relationship between white matter fractional anisotropy and other indices of cerebral health in normal aging: tract-based spatial statistics study of aging. *Neuroimage* 35, 478–487.
- Le Bihan, D., 2003. Looking into the functional architecture of the brain with diffusion MRI. *Nat. Rev. Neurosci.* 4, 469–480.
- Lemaitre, H., Crivello, F., Dufouil, C., Grassiot, B., Tzourio, C., Alperovitch, A., Mazoyer, B., 2005. No epsilon4 gene dose effect on hippocampal atrophy in a large MRI database of healthy elderly subjects. *Neuroimage* 24, 1205–1213.
- Ludwin, S.K., 1997. The pathobiology of the oligodendrocyte. *J. Neuropathol. Exp. Neurol.* 56, 111–124.
- Madden, D.J., Whiting, W.L., Huettel, S.A., White, L.E., MacFall, J.R., Provenzale, J.M., 2004. Diffusion tensor imaging of adult age differences in cerebral white matter: relation to response time. *Neuroimage* 21, 1174–1181.
- Madden, D.J., Spaniol, J., Whiting, W.L., Bucur, B., Provenzale, J.M., Cabeza, R., White, L.E., Huettel, S.A., 2007. Adult age differences in the functional neuroanatomy of visual attention: a combined fMRI and DTI study. *Neurobiol. Aging* 28, 459–476.
- Mahley, R.W., Rall Jr., S.C., 2000. Apolipoprotein E: far more than a lipid transport protein. *Annu. Rev. Genomics Hum. Genet.* 1, 507–537.
- Mahley, R.W., Weisgraber, K.H., Huang, Y., 2006. Apolipoprotein E4: a causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc. Natl Acad. Sci. USA* 103, 5644–5651.
- Mayeux, R., Small, S.A., Tang, M., Tycko, B., Stern, Y., 2001. Memory performance in healthy elderly without Alzheimer's disease: effects of time and apolipoprotein-E. *Neurobiol. Aging* 22, 683–689.
- McGeer, P.L., Schulzer, M., McGeer, E.G., 1996. Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: a review of 17 epidemiologic studies. *Neurology* 47, 425–432.
- Medina, D., Toledo-Morrell, L., Urresta, F., Gabrieli, J.D., Moseley, M., Fleischman, D., Bennett, D.A., Leurgans, S., Turner, D.A., Stebbins, G.T., 2006. White matter changes in mild cognitive impairment and AD: A diffusion tensor imaging study. *Neurobiol. Aging* 27, 663–672.
- Mosconi, L., De, S.S., Brys, M., Tsui, W.H., Pirraglia, E., Glodzik-Sobanska, L., Rich, K.E., Switalski, R., Mehta, P.D., Pratico, D., Zinkowski, R., Blennow, K., de Leon, M.J., 2008. Hypometabolism and altered cerebrospinal fluid markers in normal apolipoprotein E E4 carriers with subjective memory complaints. *Biol. Psychiatry* 63, 609–618.
- Müller, M.J., Greverus, D., Dellani, P.R., Weibrich, C., Wille, P.R., Scheurich, A., Stoeter, P., Fellgiebel, A., 2005. Functional implications of hippocampal volume and diffusivity in mild cognitive impairment. *Neuroimage* 28, 1033–1042.
- Naggara, O., Oppenheim, C., Rieu, D., Raoux, N., Rodrigo, S., Dalla, B.G., Meder, J.F., 2006. Diffusion tensor imaging in early Alzheimer's disease. *Psychiatry Res.* 146, 243–249.
- Nierenberg, J., Pomara, N., Hoptman, M.J., Sidtis, J.J., Ardekani, B.A., Lim, K.O., 2005. Abnormal white matter integrity in healthy apolipoprotein E epsilon4 carriers. *NeuroReport* 16, 1369–1372.
- Nilsson, L.G., Adolfsson, R., Backman, L., Cruts, M., Nyberg, L., Small, B.J., Van, B.C., 2006. The influence of APOE status on episodic and semantic memory: data from a population-based study. *Neuropsychology* 20, 645–657.
- Pagani, E., Agosta, F., Rocca, M.A., Caputo, D., Filippi, M., 2008. Voxel-based analysis derived from fractional anisotropy images of white matter volume changes with aging. *Neuroimage* 41, 657–667.
- Persson, J., Lind, J., Larsson, A., Ingvar, M., Cruts, M., Van, B.C., Adolfsson, R., Nilsson, L.G., Nyberg, L., 2006. Altered brain white matter integrity in healthy carriers of the APOE epsilon4 allele: a risk for AD? *Neurology* 66, 1029–1033.
- Peters, A., 2002. The effects of normal aging on myelin and nerve fibers: a review. *J. Neurocytol.* 31, 581–593.
- Peters, A., 2009. The effects of normal aging on myelinated nerve fibers in monkey central nervous system. *Front. Neuroanat.* doi:10.3389/neuro.05.011.2009.
- Pfefferbaum, A., Adalsteinsson, E., Sullivan, E.V., 2005. Frontal circuitry degradation marks healthy adult aging: evidence from diffusion tensor imaging. *Neuroimage* 26, 891–899.
- Ray, K.M., Wang, H., Chu, Y., Chen, Y.F., Bert, A., Hasso, A.N., Su, M.Y., 2006. Mild cognitive impairment: apparent diffusion coefficient in regional gray matter and white matter structures. *Radiology* 241, 197–205.
- Raz, N., Rodrigue, K.M., Kennedy, K.M., Land, S., 2009. Genetic and vascular modifiers of age-sensitive cognitive skills: effects of COMT, BDNF, ApoE, and hypertension. *Neuropsychology* 23, 105–116.
- Reiman, E.M., Caselli, R.J., Chen, K., Alexander, G.E., Bandy, D., Frost, J., 2001. Declining brain activity in cognitively normal apolipoprotein E epsilon 4 heterozygotes: a foundation for using positron emission tomography to efficiently test treatments to prevent Alzheimer's disease. *Proc. Natl Acad. Sci. USA* 98, 3334–3339.
- Reiman, E.M., Chen, K., Alexander, G.E., Caselli, R.J., Bandy, D., Osborne, D., Saunders, A. M., Hardy, J., 2004. Functional brain abnormalities in young adults at genetic risk for late-onset Alzheimer's dementia. *Proc. Natl Acad. Sci. USA* 101, 284–289.
- Reiman, E.M., Chen, K., Alexander, G.E., Caselli, R.J., Bandy, D., Osborne, D., Saunders, A. M., Hardy, J., 2005. Correlations between apolipoprotein E epsilon4 gene dose and brain-imaging measurements of regional hypometabolism. *Proc. Natl Acad. Sci. USA* 102, 8299–8302.
- Reines, S.A., Block, G.A., Morris, J.C., Liu, G., Nessly, M.L., Lines, C.R., Norman, B.A., Baranak, C.C., 2004. Rofecoxib: no effect on Alzheimer's disease in a 1-year, randomized, blinded, controlled study. *Neurology* 62, 66–71.
- Rose, S.E., Janke, A.L., Chalk, J.B., 2008. Gray and white matter changes in Alzheimer's disease: a diffusion tensor imaging study. *J. Magn. Reson. Imaging* 27, 20–26.
- Salat, D.H., Tuch, D.S., Greve, D.N., van der Kouwe, A.J., Hevelone, N.D., Zaleta, A.K., Rosen, B.R., Fischl, B., Corkin, S., Rosas, H.D., Dale, A.M., 2005. Age-related alterations in white matter microstructure measured by diffusion tensor imaging. *Neurobiol. Aging* 26, 1215–1227.
- Sando, S.B., Melquist, S., Cannon, A., Hutton, M.L., Sletvold, O., Saltvedt, I., White, L.R., Lydersen, S., Aasly, J.O., 2008. APOE epsilon 4 lowers age at onset and is a high risk factor for Alzheimer's disease; a case control study from central Norway. *BMC Neurol.* 8, 9.
- Scarmeas, N., Stern, Y., 2006. Imaging studies and APOE genotype in persons at risk for Alzheimer's disease. *Curr. Psychiatry Rep.* 8, 11–17.
- Scharf, S., Mander, A., Ugoni, A., Vajda, F., Christophidis, N., 1999. A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer's disease. *Neurology* 53, 197–201.
- Schultz, M.R., Lyons, M.J., Franz, C.E., Grant, M.D., Boake, C., Jacobson, K.C., Xian, H., Schellenberg, G.D., Eisen, S.A., Kremen, W.S., 2008. Apolipoprotein E genotype and memory in the sixth decade of life. *Neurology* 70, 1771–1777.
- Shenkin, S.D., Bastin, M.E., MacGillivray, T.J., Deary, I.J., Starr, J.M., Wardlaw, J.M., 2003. Childhood and current cognitive function in healthy 80-year-olds: a DT-MRI study. *NeuroReport* 14, 345–349.
- Shenkin, S.D., Bastin, M.E., MacGillivray, T.J., Deary, I.J., Starr, J.M., Rivers, C.S., Wardlaw, J.M., 2005. Cognitive correlates of cerebral white matter lesions and water diffusion tensor parameters in community-dwelling older people. *Cerebrovasc. Dis.* 20, 310–318.
- Small, G.W., Ercoli, L.M., Silverman, D.H., Huang, S.C., Komo, S., Bookheimer, S.Y., Lavretsky, H., Miller, K., Siddarth, P., Rasgon, N.L., Mazziotta, J.C., Saxena, S., Wu, H. M., Mega, M.S., Cummings, J.L., Saunders, A.M., Pericak-Vance, M.A., Roses, A.D., Barrio, J.R., Phelps, M.E., 2000. Cerebral metabolic and cognitive decline in persons at genetic risk for Alzheimer's disease. *Proc. Natl Acad. Sci. USA* 97, 6037–6042.
- Small, B.J., Rosnick, C.B., Fratiglioni, L., Backman, L., 2004. Apolipoprotein E and cognitive performance: a meta-analysis. *Psychol. Aging* 19, 592–600.
- Smith, C.D., Chebrolu, H., Andersen, A.H., Powell, D.A., Lovell, M.A., Xiong, S., Gold, B.T., 2008. White matter diffusion alterations in normal women at risk of Alzheimer's disease. *Neurobiol. Aging*. doi:10.1016/j.neurobiolaging.2008.08.006.
- Song, S.K., Sun, S.W., Ramsbottom, M.J., Chang, C., Russell, J., Cross, A.H., 2002. Demyelination revealed through MRI as increased radial (but unchanged axial) diffusion of water. *Neuroimage* 17, 1429–1436.
- Spreen, O., Benton, A.L., 1977. *Neurosensory Center Comprehensive Examination for Aphasia, Revised Edition*. University of Victoria Neuropsychology Laboratory, Victoria, British Columbia, Canada.
- Stahl, R., Dietrich, O., Teipel, S.J., Hampel, H., Reiser, M.F., Schoenberg, S.O., 2007. White matter damage in Alzheimer disease and mild cognitive impairment: assessment with diffusion-tensor MR imaging and parallel imaging techniques. *Radiology* 243, 483–492.
- Sullivan, E.V., Pfefferbaum, A., 2006. Diffusion tensor imaging and aging. *Neurosci. Biobehav. Rev.* 30, 749–761.
- Sullivan, E.V., Adalsteinsson, E., Pfefferbaum, A., 2006. Selective age-related degradation of anterior callosal fiber bundles quantified in vivo with fiber tracking. *Cereb. Cortex* 16, 1030–1039.
- Swan, G.E., Lessov-Schlaggar, C.N., Carmelli, D., Schellenberg, G.D., La Rue, A., 2005. Apolipoprotein E epsilon4 and change in cognitive functioning in community-dwelling older adults. *J. Geriatr. Psychiatry Neurol.* 18, 196–201.
- Sydykova, D., Stahl, R., Dietrich, O., Ewers, M., Reiser, M.F., Schoenberg, S.O., Moller, H.J., Hampel, H., Teipel, S.J., 2007. Fiber connections between the cerebral cortex and the corpus callosum in Alzheimer's disease: a diffusion tensor imaging and voxel-based morphometry study. *Cereb. Cortex* 17, 2276–2282.
- Szekely, C.A., Thorne, J.E., Zandi, P.P., Ek, M., Messias, E., Breitner, J.C., Goodman, S.N., 2004. Nonsteroidal anti-inflammatory drugs for the prevention of Alzheimer's disease: a systematic review. *Neuroepidemiology* 23, 159–169.
- Teipel, S.J., Stahl, R., Dietrich, O., Schoenberg, S.O., Pernecky, R., Bokke, A.L., Reiser, M.F., Moller, H.J., Hampel, H., 2007. Multivariate network analysis of fiber tract integrity in Alzheimer's disease. *Neuroimage* 34, 985–995.
- Walther, K., Bendlin, B.B., Glisky, E.L., Trouard, T.P., Lisse, J.R., Posever, J.O., Ryan, L., 2009. Anti-inflammatory drugs reduce age-related decreases in brain volume in cognitively normal older adults. *Neurobiol. Aging*. doi:10.1016/j.neurobiolaging.2009.03.006.
- Wechsler, D., 1981. *Wechsler Adult Intelligence Scale—Revised*. Psychological Corporation, New York.

- Wechsler, D., 1987. Wechsler Memory Scale—Revised. Psychological Corporation, New York.
- Wechsler, D., 1997. Wechsler Memory Scale—III. Psychological Corporation, San Antonio, TX.
- Wechsler, D., 1999. Wechsler Abbreviated Scale of Intelligence (WASI). Psychological Corporation, San Antonio.
- Wishart, H.A., Saykin, A.J., McAllister, T.W., Rabin, L.A., McDonald, B.C., Flashman, L.A., Roth, R.M., Mamourian, A.C., Tsongalis, G.J., Rhodes, C.H., 2006. Regional brain atrophy in cognitively intact adults with a single APOE epsilon4 allele. *Neurology* 67, 1221–1224.
- Xie, S., Xiao, J.X., Gong, G.L., Zang, Y.F., Wang, Y.H., Wu, H.K., Jiang, X.X., 2006. Voxel-based detection of white matter abnormalities in mild Alzheimer disease. *Neurology* 66, 1845–1849.
- Yasmin, H., Nakata, Y., Aoki, S., Abe, O., Sato, N., Nemoto, K., Arima, K., Furuta, N., Uno, M., Hirai, S., Masutani, Y., Ohtomo, K., 2008. Diffusion abnormalities of the uncinate fasciculus in Alzheimer's disease: diffusion tensor tract-specific analysis using a new method to measure the core of the tract. *Neuroradiology* 50, 293–299.
- Zhang, Y.T., Zhang, C.Y., Zhang, J., Li, W., 2005. Age-related changes of normal adult brain structure: analysed with diffusion tensor imaging. *Chin. Med. J. (Engl.)* 118, 1059–1065.
- Zhang, Y., Schuff, N., Jahng, G.H., Bayne, W., Mori, S., Schad, L., Mueller, S., Du, A.T., Kramer, J.H., Yaffe, K., Chui, H., Jagust, W.J., Miller, B.L., Weiner, M.W., 2007. Diffusion tensor imaging of cingulum fibers in mild cognitive impairment and Alzheimer disease. *Neurology* 68, 13–19.
- Zhang, Y., Du, A.T., Hayasaka, S., Jahng, G.H., Hlavín, J., Zhan, W., Weiner, M.W., Schuff, N., 2008. Patterns of age-related water diffusion changes in human brain by concordance and discordance analysis. *Neurobiol. Aging*. doi:10.1016/j.neurobiolaging.2008.10.009.